

Name. Parneet Singh
Student id. DE240106

Health Threats in Tourism Course
Lecturer: Mahmoud Abdou
June 20, 2025
Assignment 7



1. Explain, in no more than 500 words, how past traumatic experiences can trigger panic attacks during travel.

2. Describe, in no more than 500 words, how isolation from family and social environments during travel can lead to depression.

1. How Past Traumatic Experiences Can Trigger Panic Attacks During Travel

Traumatic experiences can leave lasting psychological imprints that alter how individuals respond to stress. When a person has previously endured trauma— such as a car accident, natural disaster, or assault— particularly in unfamiliar or mobile environments, the act of traveling can unconsciously activate those memories. The brain, especially the amygdala (responsible for processing fear), can associate similar sights, sounds, or even smells with the past event, leading to a heightened state of alertness or panic.

For example, someone who was in a serious plane crash may experience extreme anxiety or full-blown panic attacks even at the airport, due to sensory cues reminiscent of the trauma. Similarly, being in confined or crowded spaces like buses, trains, or foreign cities may replicate feelings of helplessness or entrapment from the original trauma, triggering an intense fear response. These panic attacks can involve rapid heartbeat, shortness of breath, dizziness, or the feeling of losing control— symptoms driven by the body's fight-or-flight mechanism.

Moreover, trauma can cause hypervigilance, a condition where the person is constantly on edge, scanning for danger. Traveling, with its inherent unpredictability and loss of routine, can overwhelm someone already primed for threat detection. Even minor travel-related stressors— like delays, unfamiliar customs, or language barriers— can be perceived as threats, escalating anxiety to a point where panic attacks occur.

In some cases, individuals may not consciously connect their panic with past trauma, especially if it was suppressed or never fully processed. Nonetheless, their subconscious mind continues to react defensively in certain situations. Travel thus becomes a trigger, not because of the destination itself, but because it mirrors aspects of the original traumatic experience.

Understanding this link is crucial for managing symptoms. Therapeutic techniques such as cognitive-behavioral therapy (CBT), exposure therapy, or EMDR (Eye Movement Desensitization and Reprocessing) can help retrain the brain's response and reduce the frequency or intensity of panic attacks related to travel.

2. How Isolation from Family and Social Environments During Travel Can Lead to Depression

Travel, while often associated with excitement and adventure, can also lead to emotional vulnerability, especially when it involves extended separation from familiar social and family environments. Human beings are inherently social creatures, and our emotional well-being is closely tied to our sense of connection with others. Being

removed from these support systems can leave individuals feeling adrift, lonely, and emotionally unsupported.

The absence of close relationships during travel— especially in foreign or culturally unfamiliar settings— can amplify feelings of disconnection. Without the comfort of loved ones or established routines, travelers may experience a sense of identity loss or uncertainty, particularly if they’ re dealing with major life transitions. Over time, this isolation can manifest as sadness, hopelessness, or disinterest in previously enjoyable activities— all hallmark symptoms of depression.

Language barriers, time zone differences, or cultural misunderstandings can deepen the emotional divide, making it harder to form new connections. Social withdrawal becomes both a symptom and a cause of depression, creating a cycle where the individual feels increasingly alienated and emotionally numb.

Moreover, if someone is already predisposed to depression due to genetics, previous episodes, or ongoing stress, the isolating aspects of travel can act as a catalyst. This is especially common among students studying abroad, expatriates, or professionals on long-term assignments, who may experience culture shock or homesickness that evolves into more serious depressive episodes.

Maintaining virtual contact with loved ones, establishing new social routines, and seeking local support— such as therapy or community groups— can help mitigate these effects. Recognizing the signs early and taking steps to foster connection, even in small ways, is key to preventing travel-related depression from taking root.